


Graphical Abstract

To create your abstract, type over the instructions in the template box below.
 Fonts or abstract dimensions should not be changed or altered.

γ -Selective allylic substitution reaction with Grignard reagents catalyzed by copper *N*-heterocyclic carbene complexes and its application to enantioselective synthesis

Satoshi Tominaga, Yukinao Oi, Toshio Kato,
 Duk Keun An and Sentaro Okamoto*
 Department of Applied Chemistry,
 Kanagawa University, 3-27-1 Rokkakubashi,
 Kanagawa-ku, Yokohama 221-8686, Japan



that the reaction with **4a** was slower than that with **4b** and the regioselectivity with **4a** was somewhat lower than that with **4b** (entries 1 and 2),⁸ probably due to steric bulkiness of the ligated NHC part in **4a**. The reaction could be effectively catalyzed by a reduced amount of catalyst (0.1 mol%) (entry 3). While allylic carbonate gave α -product **3** predominantly in THF (entry 4), allylic chloride afforded γ -product **2** nearly exclusively in both ether and THF (entries 7 and 8). These results suggest that coordination of the leaving group in the substrate to the metal may be important in the reaction of ester allylic substrates. On the basis of the fact that a CuCl-catalyzed reaction gave α -product exclusively (entry 9), the high γ -selectivity attained in the **4**-catalyzed reactions indicates that the NHC-copper bond (coordination) is stable and was not cleaved during the catalysis.

Table 2 summarizes representative examples of this reaction using a variety of allylic substrates and Grignard reagents. These results show the following features of this substitution reaction: (i) Primary alkyl including methyl and secondary alkyl Grignard reagents could provide **2** in excellent yield and regioselectivity. (ii) But aryl Grignard reagents gave α -product **3** predominantly (entries 4 and 5). (ii) Regarding allylic substrates, the reaction with disubstituted and trisubstituted **1** having a variety of substituents such as alkyl, aryl, siloxy- and alkoxyethyl groups with *E*- or *Z*-geometry proceeded in a highly γ -selective way.

Table 2


entry	1 (X=OCO ₂ Et)	R ⁿ MgX	2 : 3	total yield %
1	<i>n</i> -Hex	MeMgI	97 : 3	88
2	<i>n</i> -Hex	<i>n</i> -Hex.MgBr	99 : 1	quant.
3	<i>n</i> -Hex		99 : 1	quant.
4	<i>n</i> -Hex	PhMgBr ^a	2 : 98	quant.
5	<i>n</i> -Hex	PhMgBr ^a	3 : 97	82
6	Ph	<i>i</i> -PrMgCl	95 : 5	82
7	TBSO	<i>i</i> -PrMgCl	>99 : 1	quant.
8	BnO	<i>i</i> -PrMgCl	>99 : 1	quant.
9 ^b		MeMgI	> 99 : 1	84

^aPhMgBr was prepared in THF. ^b5 mol% of **4b** was used.

As shown in Scheme 1, the reaction of allylic carbonate **1** and *i*-PrMgCl with a stoichiometric amount of **4b** was found to be much slower than that with a catalytic amount of **4b** mentioned above. On this basis, the major active catalyst in the present reaction can be postulated to


be an ate-complex (higher order cuprate) such as the type [(NHC)CuR₂](MgX)⁺ but not a complex of the type (NHC)CuR⁹ although confirmation of the reaction mechanism and explanation of a high regioselectivity must await further study.

Scheme 1




With these results in hand, we turned our attention to perform the reaction in an asymmetric way by using copper complexes with chiral modified NHC ligands. Scheme 2 and Table 3 summarize preliminary results. Thus, 4-siloxy-2-buten-1-ol derivatives **1a-1d** were treated with *n*-C₆H₁₃MgBr in the presence of copper complexes with enantiomerically pure C₂-symmetric NHC ligands, **4c-4f**¹⁰ (5 mol%), in ether at -20 °C (Scheme 2). The resulting γ -product¹¹ was isolated and converted to the corresponding MTPA-esters,¹² 500 MHz ¹H NMR analyses of which confirmed the enantiomeric excess (e.e.) of the γ -product. The γ -product obtained by the reaction of (*E*)-**1d** with **4d** catalyst (entry 11) was converted to the known alcohol **5**¹³ by ozonolysis and the following reduction, and comparison of its optical rotation with that reported determined the absolute configuration as illustrated in Scheme 2.

Scheme 2



(from the reaction of (*E*)-**1d** with **4d**. See Table 3, entry 11)
 $[\alpha]_D^{23} +7.7$ (*c* 0.275, CHCl₃)
 Lit.,¹³ for *R*-isomer: $[\alpha]_D -12$ (*c* 1.0, CHCl₃).

chiral catalyst:



As can be seen from Table 3, the reactions proceeded quantitatively in S_N2' -selective fashion to give optically active γ -substituted product with low to moderate enantiomeric purity. As revealed from the results in entries 1-4, the catalyst with sterically demanding *N*-substituents (**4d**) gave the highest e.e. Interestingly, introduction of additional C_2 -chirality into the heterocyclic part of NHC resulted in inversion or decrease of the enantioselectivity (entries 3 and 4). Allylic acetate and 2-pyridyl ether¹⁴ were good substrates but carbonate and chloride gave low e.e. Inversion of the product configuration was observed when *E*-allylic substrates were used instead of *Z*-isomers, and the e.e. of the products from *E*-substrates was somewhat lower than that from the corresponding *Z*-isomers (entries 5 and 11). The highest enantiomeric ratio of 85:15 was attained by the reaction of 2-pyridyl ether (*Z*)-**1d** with catalyst **4d** (entry 10).

Table 3^a

entry	allylic substrate	catalyst	$\gamma : \alpha$	γ -product		
				% e.e. ^b	config. ^c	$[\alpha]_D^d$
1	(<i>Z</i>)- 1a	4c	87:13	40	<i>R</i>	+6.6
2		4d	95:5	60	<i>R</i>	+10.0
3		4e	88:12	38 ^e	<i>S</i>	-6.1
4		4f	84:16	6 ^e	<i>R</i>	+1.0
5	(<i>E</i>)- 1a	4d	97:3	38 ^e	<i>S</i>	-6.1
6	(<i>Z</i>)- 1b	4c	96:4	5 ^e	<i>R</i>	+0.9
7	(<i>Z</i>)- 1c	4c	77:23	8 ^e	<i>S</i>	-1.3
8		4d	84:16	16 ^e	<i>R</i>	+2.5
9	(<i>Z</i>)- 1d	4c	91:9	36	<i>R</i>	+6.3
10		4d	98:2	70	<i>R</i>	+11.0
11	(<i>E</i>)- 1d	4d	86:14	60	<i>S</i>	-9.8

^aTotal yield of the reaction was 93–100%. ^bDetermined by 500 MHz ¹H NMR analyses of the corresponding MTPA-esters, unless otherwise indicated. ^cConfirmed for entry 11 (see text). ^d c 1.0 in CHCl₃, at 23 °C. ^eBased on $[\alpha]_D$ value.

In summary, we have demonstrated that copper *N*-heterocyclic carbene (NHC) complexes are a useful catalyst for the highly S_N2' -selective allylic substitution reaction with Grignard reagents and the method can be applied to asymmetric reaction by using chiral modified NHC ligands although the preliminary results presented here reached moderate level of e.e. Further investigation on the scope of the present reaction and its synthetic utilization including development of method to attain more efficient enantioselection is underway in our laboratory.

References

- [†]Research fellow, Kanagawa University (2003). Current address; Department of Chemistry, Kangwon National University, 192-1, Hyoja2-Dong, Chunchon, Kangwon-Do, 200-701 Korea.
- Lipshutz, B. H.; Senguota, S. *Org. React.*, **1992**, *41*, 135.
- For organocopper chemistry, see: “*Organocopper Reagents*,” R. J. K. Talor Ed., Oxford University Press, Oxford (1994).
- Lipshutz, B. H. in “*Organometallics in Synthesis*,” M. Schlosser Ed., Wiley, Chichester (2002), p. 665. “*Modern Organocopper Chemistry*,” N. Krause Ed., Wiley-VHC, Weinheim (2001).
- Chiral copper complex-catalyzed enantioselective allylic substitution reactions: with Grignard reagents: van Klaveren, M.; Persson, E. S. M.; del Villar, A.; Grove, D. M.; Bäckvall, J.-E.; van Koten, G. *Tetrahedron Lett.*, **1995**, *36*, 3059. Karlström, A. S. E.; Huerta, F. F.; Meuzelaar, G. J.; Bäckvall, J.-E. *Synlett*, **2001**, 923. Meuzelaar, G. J.; Karlström, A. S. E.; van Klaveren, M.; Persson, E. S. M.; del Villar, S.; van Koten, G.; Bäckvall, J.-E. *Tetrahedron*, **2000**, *56*, 2895. Alexakis, A.; Malan, C.; Lea, L.; Benhaim, C.; Fournieux, X. *Synlett*, **2001**, 927. Alexakis, A.; Croset, K. *Org. Lett.*, **2002**, *4*, 4147. Croset, K.; Polet, D.; Alexakis, A. *Angew. Chem., Int. Ed. Engl.*, **2004**, *43*, 2426. With dialkylzinc compounds, Dübner, F.; Knochel, P. *Angew. Chem., Int. Ed. Engl.*, **1999**, *38*, 379. Dübner, F.; Knochel, P. *Tetrahedron Lett.*, **2000**, *41*, 9233. Malda, H.; van Zijl, A. W.; Arnold, L. A.; Feringa, B. L. *Org. Lett.*, **2001**, *3*, 1169. Luchaco-Cullis, C. A.; Mizutani, H.; Muphy, K. E.; Hoveyda, A. H. *Angew. Chem., Int. Ed. Engl.*, **2001**, *40*, 1456.
- Asymmetric conjugate addition reactions catalyzed by copper imidazolium carbene complexes: (a) Guillen, F.; Winn, C. L.; Alexakis, A. *Tetrahedron: Asymmetry*, **2001**, *12*, 2083. (b) Fraser, P. K.; Woodward, S. *Tetrahedron Lett.*, **2001**, 42, 2747. (c) Alexakis, A.; Winn, C. L.; Guillen, F.; Pytkowicz, J.; Roland, S.; Mangeney, P. *Adv. Synth. Catal.*, **2003**, *345*, 345.
- Other catalyses with copper imidazolium carbene complexes in organic synthesis: (a) Jurkauskas, V.; Sadighi, J. P.; Buchwald, S. L. *Org. Lett.*, **2003**, *5*, 2417. (b) Kaur, H.; Zinn, F. K.; Stevens, E. D.; Nolan, S. P. *Organometallics*, **2004**, *23*, 1157 and references for recent structural studies of *N*-heterocyclic carbene copper complexes cited therein.
- Compounds **4a**⁵ and **4b** were prepared by the reaction of CuCl, *t*-BuONa and the corresponding imidazolium chloride in THF according to the reported procedure.⁵ The isolated **4a** and **4b** were used. **4b**: Mp. 268 °C; ¹H NMR (CDCl₃, 500 MHz) δ 7.31 (t, *J* = 7.5 Hz, 2H), 7.21 (d, *J* = 8.0 Hz, 4H), 7.10 (s, 2H), 2.17 (s, 12H); ¹³C NMR (CDCl₃, 125 MHz) δ 137.4, 134.9, 129.6, 128.8, 122.1, 17.8 (one carbon could not be detected); IR (neat) 3166, 1695, 1599, 1557, 1479, 1443, 1404, 1335, 1290, 1230, 1161, 1110, 1038, 948, 786, 735, 693, 609 cm⁻¹. Elemental analysis, Calcd. for C₁₉H₂₂N₂OClCu; C, 60.79; H, 5.37; N, 7.46. Found; C, 60.95; H, 5.54; N, 7.47.
- To a suspension of CuCl-NHC complex **4** (0.02 mmol) in ether (1 mL) was added Grignard reagent (0.7–1.5 M in ether, 3.0 mmol) at 0 °C and the mixture was stirred for 0.5 h at this temperature. To this was added a solution of allylic substrate **1** (2.0 mmol) in ether (2 mL) and the mixture was stirred at 0 °C. After confirming the completion of the reaction by TLC analysis, saturated aqueous NH₄Cl was added. Usual extractive work-up and passing through a short silica gel column yielded a mixture of α - and γ -products.
- TLC analysis of the reaction mixture indicated that for complete consumption of **1a** it needed 2 h for the reaction with **4a** and 30min for the reaction with **4b**.
- Synthesis and structural study of (NHC)Cu(I) methyl complex were reported: Mankad, N. P.; Gray, T. G.; Laitar, D. S.; Sadighi, J. P. *Organometallics*, **2004**, *23*, 1191.
- Compounds **4c–4f** were prepared from the corresponding imidazolium salts,^{4c} *t*-BuONa and CuCl in THF. The resulting mixture was filtered through a pad of Celite and the filtrate was stored under Ar. The THF-solution containing **4c–4f**, respectively, thus obtained was charged into the reaction

vessel and THF was removed under reduced pressure prior to use.

11. ***tert*-Butyldimethyl(2-hexylbut-3-enyloxy)silane**: ^1H NMR (CDCl_3 , 270 MHz) δ 5.62 (ddd, $J = 8.4, 9.7, 17.8$ Hz, 1H), 4.97-5.07 (m, 2H), 3.52 (dd, $J = 6.3, 11.2$ Hz, 1H), 3.49 (dd, $J = 6.3, 11.2$ Hz, 1H), 2.07-2.23 (m, 1H), 1.10-1.60 (m, 10H), 0.89 (s, 9H), 0.88 (t, $J = 7.1$ Hz, 3H), 0.07 and 0.04 (2s, each 3H); ^{13}C NMR (CDCl_3 , 67.5 MHz) δ 140.6, 115.2, 66.7, 31.8, 30.8, 29.4, 27.0, 26.0, 22.7, 18.4, 14.1, -5.3, -5.4; IR (neat) 3076, 2926, 2854, 1644, 1470, 1383, 1365, 1254, 1101, 1008, 942, 915, 837, 777 cm^{-1} . Elemental analysis, Calcd. for $\text{C}_{16}\text{H}_{34}\text{OSi}$; C, 71.04; H, 12.67. Found; C, 70.64; H, 13.00.
12. Dale, J. A.; Mosher, H. S. *J. Am. Chem. Soc.*, **1971**, *95*, 512.
13. Gurjar, M. K.; Bhaket, P. *Heterocycles*, **2000**, *53*, 143.
14. We found that the reaction of allylic 2-pyridyl ethers with alkyl Grignard reagents in the presence of a catalytic amount of CuCN in ether proceeded highly predominantly in an $\text{S}_{\text{N}}2'$ fashion (unpublished results).